

Unclassified

AD-A258 551

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for Public Release Distribution Unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. DU/DC/TR-32			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Department of Chemistry Duke University		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research		
6c. ADDRESS (City, State, and ZIP Code) Durham, NC 27708-0346			7b. ADDRESS (City, State, and ZIP Code) 300 North Quincy Street Arlington, VA 22217-5000		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-89-J1545 R&T Code 4135008		
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. NR	PROJECT NO. 053	TASK NO. 841
11. TITLE (Include Security Classification) X-RAY CRYSTAL STRUCTURE OF $\text{InCl}_3 \cdot 2\text{THF}$ , A NEUTRAL FIVE-COORDINATE BIS-ADDUCT OF AN IN(III) TRIHALIDE					
12. PERSONAL AUTHOR(S) Mark F. Self, Andrew T. McPhail, and Richard L. Wells*					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 1992-11-18	
15. PAGE COUNT 12					
16. SUPPLEMENTARY NOTATION Accepted for Publication in <i>Polyhedron</i>					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) indium trichloride, adduct, five-coordinate, synthesis, crystal structure		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  See Attached  118225 92-30111 bpr					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Richard L. Wells			22b. TELEPHONE (Include Area Code) (919) 660-1541		22c. OFFICE SYMBOL

[19. ABSTRACT]

The 2:1 mole ratio reaction of  $\text{InCl}_3$  and  $\text{P}(\text{SiMe}_3)_3$  in a toluene/THF mixture was investigated. Reaction appears to occur in a 1:1 mole fashion, since the products obtained are unreacted  $\text{InCl}_3$  (48.5% recovered) as  $\text{InCl}_3 \cdot 2\text{THF}$  and a fine powder which roughly analyzes to be  $[\text{Cl}_2\text{In}_3\text{P}_3(\text{SiMe}_3)_2]_n$ . The latter was characterized by melting point and partial elemental analysis. The X-ray crystal structure of  $\text{InCl}_3 \cdot 2\text{THF}$  is reported.  $\text{InCl}_3 \cdot 2\text{THF}$  crystallizes in the orthorhombic space group  $Pbcn$  [ $(D_{2h}^{14})$ , No.60] with unit cell parameters of  $a = 10.802(1) \text{ \AA}$ ,  $b = 10.101(1) \text{ \AA}$ ,  $c = 13.083(1) \text{ \AA}$ ,  $V = 1427.5(4) \text{ \AA}^3$ , and  $D_{\text{calcd}} = 1.700 \text{ g cm}^{-3}$  for  $Z = 4$  [ $R = 0.042$  ( $R_w = 0.058$ )] and is an example of a neutral pentacoordinated bis-adduct of an indium(III) trihalide. The crystal structure consists of discrete monomeric units with no abnormally short intermolecular separations. The geometry about the indium atom approximates to trigonal bipyramidal with oxygen atoms of each of the THF molecules occupying axial positions [ $\text{In-O} = 2.257(9) \text{ \AA}$ ,  $\text{O-In-O} = 178.5(3)^\circ$ ] while the three chlorine atoms (mean  $\text{In-Cl} = 2.321 \text{ \AA}$ ) lie in the equatorial plane.

DTIC QUALITY INSPECTED 4

Accession For		
NTIS	CRA&I	<input checked="" type="checkbox"/>
DTIC	TAB	<input type="checkbox"/>
Unannounced		<input type="checkbox"/>
Justification		
By		
Distribution /		
Availability Codes		
Dist	Avail and/or Special	
A-1		

OFFICE OF NAVAL RESEARCH

Grant NO0014-89-J-1545

R&T Code 4135008

Technical Report No. DU/DC/TR-32

X-RAY CRYSTAL STRUCTURE OF  $\text{InCl}_3 \cdot 2\text{THF}$ ,  
A NEUTRAL FIVE-COORDINATE BIS-ADDUCT OF AN  $\text{In(III)}$  TRIHALIDE

by

Mark F. Self, Andrew T. McPhail, and Richard L. Wells\*

Prepared for Publication in *Polyhedron*

Duke University  
Department of Chemistry,  
P. M. Gross Chemical Laboratory  
Box 90346  
Durham, NC 27708-0346

November 18, 1992

Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

**X-Ray Crystal Structure of  $\text{InCl}_3 \cdot 2\text{THF}$ , a Neutral Five-Coordinate  
Bis-adduct of an In(III) Trihalide**

**Mark F. Self, Andrew T. McPhail, and Richard L. Wells\***

Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke  
University, Durham, North Carolina 27708, U.S.A.

(Received September 14, 1992; accepted ....., 1992)

**Abstract** - The 2:1 mole ratio reaction of  $\text{InCl}_3$  and  $\text{P}(\text{SiMe}_3)_3$  in a toluene/THF mixture was investigated. Reaction appears to occur in a 1:1 mole fashion, since the products obtained are unreacted  $\text{InCl}_3$  (48.5% recovered) as  $\text{InCl}_3 \cdot 2\text{THF}$  and a fine powder which roughly analyzes to be  $[\text{Cl}_2\text{In}_3\text{P}_3(\text{SiMe}_3)_2]_n$ . The latter was characterized by melting point and partial elemental analysis. The X-ray crystal structure of  $\text{InCl}_3 \cdot 2\text{THF}$  is reported.  $\text{InCl}_3 \cdot 2\text{THF}$  crystallizes in the orthorhombic space group  $Pbcn$  [ $(D_{2h}^{14})$ , No.60] with unit cell parameters of  $a = 10.802(1) \text{ \AA}$ ,  $b = 10.101(1) \text{ \AA}$ ,  $c = 13.083(1) \text{ \AA}$ ,  $V = 1427.5(4) \text{ \AA}^3$ , and  $D_{\text{calcd}} = 1.700 \text{ g cm}^{-3}$  for  $Z = 4$  [ $R = 0.042$  ( $R_w = 0.058$ )] and is an example of a neutral pentacoordinated bis-adduct of an indium(III) trihalide. The crystal structure consists of discrete monomeric units with no abnormally short intermolecular separations. The geometry about the indium atom approximates to trigonal bipyramidal with oxygen atoms of each of the THF molecules occupying axial positions [ $\text{In-O} = 2.257(9) \text{ \AA}$ ,  $\text{O-In-O} = 178.5(3)^\circ$ ] while the three chlorine atoms (mean  $\text{In-Cl} = 2.321 \text{ \AA}$ ) lie in the equatorial plane.

---

\* Author to whom correspondence should be addressed.

For some time our laboratory has been investigating convenient and facile routes to the formation of novel compounds containing Ga-As bonds.<sup>1</sup> In fact, one such compound with the empirical formula  $(\text{AsCl}_3\text{Ga}_2)_n$  (**I**) can be easily converted into nanocrystalline GaAs itself.<sup>2</sup> Compound **I** results from the 2:1 mole ratio reaction of  $\text{GaCl}_3$  and  $\text{As}(\text{SiMe}_3)_3$ . In related work, Barron and coworkers recently reported that the 1:1 mole ratio reaction of  $\text{InCl}_3$  and  $\text{P}(\text{SiMe}_3)_3$  at low temperature yields  $[\text{Cl}_2\text{InP}(\text{SiMe}_3)_2]_x$  (**II**).<sup>3</sup> In that we have now begun to direct considerable attention to the production of compounds containing In-P bonds,<sup>4-7</sup> it was decided that the In/P analogue of the reaction leading to the formation of **I** should be investigated [i.e., the 2:1 reaction of  $\text{InCl}_3$  and  $\text{P}(\text{SiMe}_3)_3$ ], the goal being the possible isolation of new precursor  $(\text{PCl}_3\text{In}_2)_n$ . We report that though the 2:1 reaction of  $\text{GaCl}_3$  and  $\text{As}(\text{SiMe}_3)_3$  to produce **I** occurs cleanly and readily, the corresponding reaction of  $\text{InCl}_3$  and  $\text{P}(\text{SiMe}_3)_3$  affords  $[\text{Cl}_2\text{In}_3\text{P}_3(\text{SiMe}_3)_2]_n$  (**III**) and unreacted  $\text{InCl}_3$  (48.5% of the starting  $\text{InCl}_3$ ) in a solvated form  $[\text{InCl}_3 \cdot 2\text{THF}]$  (**IV**). Compound **III** (characterized by melting point and partial elemental analysis) was isolated as a relatively unreactive fine orange powder. It is insoluble in hydrocarbon solvents, THF, and  $\text{Et}_2\text{O}$ , and air stable for several hours at room temperature. Complex **IV** has been previously synthesized and quite extensively characterized by Fairbrother *et. al.*,<sup>8</sup> but no solid-state structure was ever obtained. Herein, we report the crystal structure of **IV** as it represents a novel example of a neutral pentacoordinated bis-adduct of an indium(III) trihalide.

## EXPERIMENTAL

### *Synthesis*

All manipulations were performed using general Schlenk techniques. Colorless crystals of **IV** were obtained by the 2:1 mole reaction of  $\text{P}(\text{SiMe}_3)_3$ <sup>9</sup> and  $\text{InCl}_3$  (purchased from Strem chemicals). In the dry box (Vacuum/Atmospheres HE-493 Dri-Lab containing an argon atmosphere),  $\text{P}(\text{SiMe}_3)_3$  (0.250 g, 0.998 mmol) was dissolved in 25 cm<sup>3</sup> of anhydrous, degassed toluene (distilled from sodium/benzophenone ketyl) in a 100-cm<sup>3</sup> one-necked round-bottomed flask equipped with a Teflon valve and a micro-stirbar. A suspension of  $\text{InCl}_3$  (0.441 g, 1.99 mmol) in 15 cm<sup>3</sup> of toluene and 15 cm<sup>3</sup> of anhydrous, degassed tetrahydrofuran (THF) (distilled from sodium/benzophenone ketyl) was slowly added drop-wise. Immediately upon addition, the reaction mixture became yellow-orange and quite turbid. The reaction mixture was stirred at room temperature for 2 days then allowed to settle undisturbed for 1 day. Decantation of the colorless mother liquor from the bright orange powder  $\{[\text{Cl}_2\text{In}_3\text{P}_3(\text{SiMe}_3)_2]_n$  (**III**), 0.204g, 47% yield based on In, Mp = 200 to 260°(dec to black), Anal. Calcd (Found) for  $\text{C}_6\text{H}_{18}\text{Cl}_2\text{In}_3\text{P}_3\text{Si}_2$ : C, 11.61 (11.40, 11.62); H, 2.75 (2.59, 2.53); Cl, 10.83 (10.66, 10.91) and subsequent cooling at -15° C for 3 days afforded crystals of **IV** (0.352g, 48.5% yield based on In) suitable for X-ray analysis.

### *X-ray structural solution and refinement*

Crystallographic data are summarized in Table 1. The crystal used was a colorless block mounted inside a flame-sealed 0.6 mm thin-walled glass capillary under an inert argon atmosphere. Oscillation and Weissenberg photographs yielded preliminary unit cell parameters and

space group information. X-ray intensity data were recorded on an Enraf-Nonius CAD-4 diffractometer at 25° C using graphite-monochromated Cu-K $\alpha$  radiation ( $\lambda = 1.5418 \text{ \AA}$ ). Lorentz and polarization corrections were applied as was an empirical absorption correction. The crystal structure was solved by direct methods. Initial coordinates for the indium and chlorine atoms were derived from an *E*-map. Carbon and oxygen atoms were located in a difference Fourier synthesis phased by these atoms. Several rounds of full-matrix least-squares adjustment of non-hydrogen atom positional and thermal parameters (at first isotropic, then anisotropic) followed. Hydrogen atoms were incorporated at their calculated positions (C-H = 1.05  $\text{\AA}$ ) in the later iterations which converged (max. shift/esd = 0.02) at  $R = 0.042$  ( $R_w = 0.058$ ). A final difference Fourier synthesis revealed no unusual features (max. 0.56, min. -0.41  $\text{e/\AA}^3$ , both in the vicinity of the indium atom). Crystallographic calculations were performed using the Enraf-Nonius Structure Determination Package (SDP).<sup>10</sup> For all structure-factor calculations, neutral atom scattering factors and their anomalous dispersion corrections were taken from ref. 11. Selected distances and angles are provided in Table 2. An ORTEP diagram showing the atom labeling scheme and solid-state conformation of **IV** is presented in Figure 1; the packing arrangement of molecules of **IV** in the crystal is illustrated in Figure 2. Full information concerning conditions for crystallographic data collection and structure refinement, atomic coordinates, positional parameters, anisotropic thermal parameters, and observed and calculated structure factors have been deposited with the Cambridge Crystallographic Data Centre.

## RESULTS AND DISCUSSION

Though the structural chemistry of five-coordinate aluminum and gallium compounds has been well examined, corresponding investigations of such indium compounds have remained more sparse.<sup>12</sup> Currently, there are seven pentacoordinate indium(III) trihalide bis-adducts which have been crystallographically characterized [*viz.*,  $\text{InX}_3 \cdot \text{L}_2$ : X = Cl; L =  $\text{PPh}_3$  (V),<sup>13</sup> HMPA ( $\text{OP}\{\text{N}(\text{Me})_2\}_3$ ) (VI),<sup>14</sup>  $\text{NMe}_3$  (VII),<sup>15</sup>  $\text{SPMe}_3$  (VIII),<sup>16</sup>  $\text{SAsMe}_3$  (IX)<sup>16</sup>; X = Br; L =  $\text{SPMe}_3$  (X),<sup>16</sup>  $\text{SAsMe}_3$  (XI)<sup>16</sup>]. Of these, it is interesting to note that five (VII - XI) were only reported within the last six years and IV is only the second such compound containing In-O bonds to be structurally characterized, the first being VI.

The crystal structure of IV consists of discrete monomeric units which have one of their In-Cl bonds lying on a crystallographic  $C_2$  symmetry axis (see Fig. 1). In common with the other  $\text{InX}_3 \cdot \text{L}_2$  adducts, the coordination geometry about the indium atoms is trigonal-bipyramidal with three chlorine atoms defining the equatorial plane and the oxygen atoms of the THF molecules occupying the axial sites. The O-In-O angle at  $178.5(3)^\circ$  in IV deviates only slightly from an exactly linear value and lies close to that of  $178.0(2)^\circ$  in VI. The In-O bond length of  $2.257(9) \text{ \AA}$  in IV is significantly longer and thus weaker than the corresponding bonds in VI where they average  $2.180 \text{ \AA}$ ; this difference may be ascribed to the increased electron donating ability of the oxygen atom in the HMPA ligand. The mean In-Cl bond length at  $2.321 \text{ \AA}$  in IV is shorter than that of  $2.359 \text{ \AA}$  in VI as well as the corresponding lengths of  $2.360 \text{ \AA}$  -  $2.412 \text{ \AA}$  in V, VII - IX. The observed decrease in the equatorial bond lengths in IV vs. VI is correlated with the aforementioned lengthened axial bond distance in the



former. Endocyclic torsion angles characterizing the conformation of the THF molecule are related by an approximate  $C_2$  symmetry axis which passes through the oxygen atom and the midpoint of the C(3)-C(4) bond, and, accordingly, this ring is in a half-chair conformation. The mean bond angle subtended at the oxygen atom is  $118^\circ$  and thus the geometry about this atom lies closer to trigonal planar than to tetrahedral. X-ray crystallographic studies<sup>16</sup> on the series  $\text{InX}_3 \cdot \text{L}_2$  ( $\text{X} = \text{Cl}, \text{Br}$ ;  $\text{L} = \text{SPMe}_3, \text{SAsMe}_3$ ) have shown that in the solid state the angle subtended at the axial sulfur atom is dependent on intermolecular as well as intramolecular non-bonded interactions. The orientation of the In-O bond with respect to the THF ring in crystals of IV produces no unusually short intramolecular Cl...H non-bonded distances (min. Cl...H = 3.29 Å). Crystal packing requirements involving two equidistant intermolecular interactions between Cl(2) in the reference molecule and one of the hydrogen atoms at C(2) in neighboring molecules [Cl(2)...H(2) = 2.87 Å] apparently dictate the geometry at the oxygen atom in the solid state.

*Acknowledgement* - This work was financially supported by the Office of Naval Research.

## References

- 1) For a general review see R. L. Wells, *Coord. Chem. Rev.* 1992, **112**, 273.
- 2) R. L. Wells, R. B. Hallock, A. T. McPhail, C. G. Pitt, and J. D. Johansen, *Chem. Mater.* 1991, **3**, 381.
- 3) M. D. Healy, P. E. Laibinis, P. D. Stupik, and A. R. Barron, *J. Chem. Soc., Chem. Commun.* 1989, 359.
- 4) R. L. Wells, A. T. McPhail, and M. F. Self, *Organometallics* 1992, **11**, 221.
- 5) R. L. Wells, A. T. McPhail, L. J. Jones, M. F. Self, and R. J. Butcher, *Organometallics* 1992, **11**, 2694.
- 6) R. L. Wells, A. T. McPhail, and M. F. Self, *J. Coord. Chem.* **in press**.
- 7) R. L. Wells, A. T. McPhail, L. J. Jones, and M. F. Self, *J. Organomet. Chem.* **in press**.
- 8) F. Fairbrother, N. Flitcroft, and H. Prophet, *J. Less-Common Metals* 1960, **2**, 49.
- 9) G. Becker, and W. Holderich, *Chem. Ber.* 1975, **108**, 2484.
- 10) Enraf-Nonius Structure Determination Package (SDP 3.0), Enraf-Nonius, Delft, The Netherlands, 1985.
- 11) *International Tables for X-ray Crystallography*; The Kynoch Press, Birmingham, England, 1974, Vol IV.
- 12) (a) D. G. Tuck, In *Comprehensive Organometallic Chemistry*; G. Wilkinson, Ed; Pergamon Press: Oxford, England, 1982; Vol. 1, Chpt. 7, pp. 683 - 723. (b) D. G. Tuck, In *Comprehensive Coordination Chemistry*; G. Wilkinson, R. D. Gillard, and J. A. McCleverty, Eds; Pergamon Press: Oxford, England, 1987; Vol. 3, Chpt. 25.2, pp. 153 - 182.
- 13) M. V. Veidis, and G. J. Palenik, *J. Chem. Soc. D* 1969, 586.
- 14) S. P. Sinha, T. T. Pakkanen, T. A. Pakkanen, and L. Niinisto, *Polyhedron* 1982, **1**, 355.
- 15) R. Karia, G. R. Willey, and M. G. B. Drew, *Acta. Cryst. Sect. C. Cryst. Struct. Comm.* 1986, **C42**, 558.
- 16) W. T. Robinson, C. J. Wilkins, and Z. Zeying, *J. Chem. Soc., Dalton Trans.* 1988, 2187.

Figure captions

Fig. 1. ORTEP diagram (50% probability ellipsoids) showing the atom numbering scheme and solid-state conformation of  $\text{InCl}_3 \cdot 2\text{THF}$  (IV); primed atoms are related to the unprimed atoms by a crystallographic  $C_2$  symmetry axis passing through In and Cl(2). Small circles represent hydrogen atoms.

Fig. 2. Packing arrangement of molecules of  $\text{InCl}_3 \cdot 2\text{THF}$  (IV) in the crystal.




Fig. 1




Fig. 2

Table 1. Crystallographic data for  $\text{InCl}_3 \cdot 2\text{THF}$  (IV)

molecular formula	$\text{C}_8\text{H}_{16}\text{Cl}_3\text{InO}_2$
formula weight	365.39
crystal system	orthorhombic
space group	$Pbcn(D_{2h}^{14})$ - No.60
$a(\text{\AA})$	10.802(1)
$b(\text{\AA})$	10.101(1)
$c(\text{\AA})$	13.083(1)
no. of orientation refls.; $\theta(^{\circ})$ range	25; 36-40
$V(\text{\AA}^3)$	1427.5(4)
$Z$	4
$D_{\text{calcd.}} (\text{g cm}^{-3})$	1.700
$\mu(\text{Cu-K}\alpha \text{ radiation}, \lambda = 1.5418 \text{ \AA})$	186.9
temp. ( $^{\circ}\text{C}$ )	25
crystal dimensions (mm)	0.16 x 0.18 x 0.20
$T_{\text{max}}:T_{\text{min}}$	1.00:0.71
scan type	$\omega$ -2 $\theta$
scanwidth ( $^{\circ}$ )	$0.80 + 0.14\tan\theta$
$\theta_{\text{max}}(^{\circ})$	75
intensity control refls.;	1 1 1, 1 2 2, 1 3 3, 1 1 4
variation; repeat time (hr)	<1% ; 2
total no. of refls. ( $+h, +k, \pm l$ ) recorded	1471
no. of refls. retained [ $I > 3.0\sigma(I)$ ]	537
no. of parameters refined	65
$R(R_w)^a$	0.042 (0.058)
goodness-of-fit <sup>b</sup>	1.16
max. shift:esd in final least-squares cycle	0.02
final $\Delta\rho(\text{e/\AA}^3)$ max.;min.	0.56 ; -0.41

<sup>a</sup> $R = \Sigma ||F_o| - |F_c|| / \Sigma |F_o|$ ;  $R_w = [\Sigma w(|F_o| - |F_c|)^2 / \Sigma w |F_o|^2]^{1/2}$ ;  $\Sigma w \Delta^2$  [ $w = 1/\sigma^2(|F_o|)$ ,  $\Delta = (|F_o| - |F_c|)$ ] was minimized.

<sup>b</sup>Goodness-of-fit =  $[\Sigma w \Delta^2 / (N_{\text{observations}} - N_{\text{parameters}})]^{1/2}$ .

Table 2. Interatomic Distances (Å) and Angles<sup>a</sup> (deg.) for InCl<sub>3</sub>·2THF (IV), with ESD's in Parentheses

---

a) Bond Distances

In-Cl(1)	2.330(4)	O(1)-C(5)	1.49(2)
In-Cl(2)	2.312(5)	C(2)-C(3)	1.53(3)
In-O(1)	2.257(9)	C(3)-C(4)	1.45(3)
O(1)-C(2)	1.46(2)	C(4)-C(5)	1.50(2)

(b) Bond Angles

Cl(1)-In-Cl(1')	117.8(2)	In-O(1)-C(2)	122(1)
Cl(1)-In-Cl(2)	121.1(1)	In-O(1)-C(5)	122(1)
Cl(1)-In-O(1)	90.5(3)	C(2)-O(1)-C(5)	109(1)
Cl(1)-In-O(1')	90.3(3)	O(1)-C(2)-C(3)	104(1)
Cl(2)-In-O(1)	89.2(2)	C(2)-C(3)-C(4)	105(1)
O(1)-In-O(1')	178.5(3)	C(3)-C(4)-C(5)	104(1)
		O(1)-C(5)-C(4)	104(1)

---

<sup>a</sup>Primed atoms are related to unprimed atoms by the following transformation:  
 $-x, y, 1/2 - z$ .

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (2)\*  
Chemistry Division, Code 1113  
800 North Quincy Street  
Arlington, Virginia 22217-5000

Dr. James S. Murday (1)  
Chemistry Division, Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Robert Green, Director (1)  
Chemistry Division, Code 385  
Naval Air Weapons Center  
Weapons Division  
China Lake, CA 93555-6001

Dr. Elek Lindner (1)  
Naval Command, Control and Ocean  
Surveillance Center  
RDT&E Division  
San Diego, CA 92152-5000

Dr. Bernard E. Douda (1)  
Crane Division  
Naval Surface Warfare Center  
Crane, Indiana 47522-5000

Dr. Richard W. Drisko (1)  
Naval Civil Engineering  
Laboratory  
Code L52  
Port Hueneme, CA 93043

Dr. Harold H. Singerman (1)  
Naval Surface Warfare Center  
Carderock Division Detachment  
Annapolis, MD 21402-1198

Dr. Eugene C. Fischer (1)  
Code 2840  
Naval Surface Warfare Center  
Carderock Division Detachment  
Annapolis, MD 21402-1198

Defense Technical Information  
Center (2)  
Building 5, Cameron Station  
Alexandria, VA 22314